

5 SEM TDC PHY M 1

2 0 1 6

(November)

PHYSICS

(Major)

Course : 501

(**Mathematical Physics**)

Full Marks : 60

Pass Marks : 24 (Backlog)/18 (2014 onwards)

Time : 3 hours

The figures in the margin indicate full marks for the questions

1. Choose the correct option : 1×6=6

(a) Which of the following integrals is non-vanishing?

(i) $\int_{-1}^{+1} x P_n \frac{dP_m}{dx} dx$ for $n > m$

(ii) $\int_{-1}^{+1} P_n(x) dx$

(iii) $\int_{-1}^{+1} x^2 P_5(x) dx$

(iv) $\int_{-1}^{+1} P_0(x) dx$

(b) Given $\Gamma(3)\Gamma\left(\frac{5}{2}\right) = A\Gamma(6)$, find A.

(i) $\sqrt{\pi}$

(ii) $\sqrt{\pi}/2$

(iii) $\sqrt{\pi}/2^3$

(iv) $\sqrt{\pi}/2^5$

(c) If $u = x^3 - 3xy^2$, the analytic function $f(z) = u + iv$ will be

(i) z^3

(ii) z^{-3}

(iii) $|z|^3$

(iv) None of the above

(d) What is the ratio of coefficients of z^n and $\frac{1}{z^n}$ in the Laurent's expansion of

the function $\cosh\left(z + \frac{1}{z}\right)$?

(i) 0

(ii) $\frac{1}{2}$

(iii) 1

(iv) None of the above

(e) The value of a_0 in the Fourier series of t^2 in the interval $-\pi < t < \pi$ is

(i) 0

(ii) $\frac{\pi^2}{3}$

(iii) $\frac{\pi^2}{8}$

(iv) $\frac{\pi^2}{4}$

(f) Using Fourier integral, the value of $\int_0^{\infty} \frac{\cos xu}{1+u^2} du$ ($x > 0$) is found to be

(i) $\frac{\pi}{2}$

(ii) $\frac{\pi}{2} e^x$

(iii) $\frac{2}{\pi} e^{-x}$

(iv) $\frac{\pi}{2} e^{-x}$

2. (a) Prove that

$$\sum_{n=1}^{\infty} \frac{1}{(2n-1)^2} = \frac{\pi^2}{8}$$

(b) Find the value of a_n in the Fourier series of $f(x)$ in the interval $(-\pi, \pi)$, where

$$\begin{aligned} f(x) &= \pi + x, \text{ when } -\pi < x < 0 \\ &= \pi - x, \text{ when } 0 < x < \pi \end{aligned}$$

2

(c) Prove that $P_{2m}(-\mu) = P_{2m}(\mu)$.

2

(d) Express the integral $I = \int_0^{\infty} \frac{x^3}{(1+x)^5} dx$ in terms of beta and gamma functions and hence find its value.

2

(e) Using Cauchy's integral formula, evaluate the integral $\oint \frac{z^2}{(z^2-1)} dz$ around the unit circle with centre at $z=1$.

2

(f) If $u(x, y) = x^2 - y^2$ is the real part of an analytic function $f(z) = u + iv$, find v .

2

3. (a) Solve the equation $y'' - y = 0$ with $y(0) = 4, y'(0) = -2$.

3

(b) Find the solution of the non-homogeneous equation $y'' + 4y = 8x^2$.

3

(c) Prove that

$$P_n(x) = \frac{1}{2^n n!} \left(\frac{d}{dx} \right)^n (x^2 - 1)^n$$

4

(d) Prove that

$$(2n+1)xP_n(x) = (n+1)P_{n+1}(x) + nP_{n-1}(x) \quad 4$$

Or

Prove that Legendre polynomial $P_n(\mu)$ is the coefficient of h^2 in $(1 - 2\mu h + h^2)^{-1/2}$. 4

4. (a) Prove that if $f(z) = u(x, y) + iv(x, y)$ is analytic in a domain D , then u and v satisfy $\nabla^2 u = 0$ and $\nabla^2 v = 0$. 4

(b) Prove that if $f(z)$ is an analytic function on and within the closed contour c , the value of $f(z)$ at a point $z = \epsilon$ inside c is given by

$$f(\epsilon) = \frac{1}{2\pi i} \oint \frac{f(z)}{z - \epsilon} dz \quad 4$$

(c) Answer any two from the following :

3×2=6

(i) Show that the triangle whose vertices are the points z_1, z_2, z_3 in Argand diagram will be equilateral if

$$z_1^2 + z_2^2 + z_3^2 = z_1 z_2 + z_2 z_3 + z_3 z_1$$

(ii) If $f(z)$ is an analytic function of $|z|$, prove that

$$\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) |f(z)|^2 = 4 |f'(z)|^2$$

(iii) Obtain the expansion

$$f(z) = f(a) + 2 \left\{ \frac{z-a}{2} f' \left(\frac{z+a}{2} \right) + \frac{(z-a)^3}{2^3 3!} f''' \left(\frac{z+a}{2} \right) + \frac{(z-a)^5}{2^5 5!} f^{(5)} \left(\frac{z+a}{2} \right) + \dots \right\}$$

and determine its range of validity.

5. (a) Find an even function of x which is equal to kx for $0 \leq x \leq l/2$ and is

$$k(l-x) \text{ for } l/2 \leq x \leq l$$

3

(b) Find the series of sines and cosines of multiples of x which represents $f(x)$ in the interval $-\pi < x < \pi$, where

$$f(x) = 0, \text{ when } -\pi < x < 0$$

$$= \frac{\pi x}{4}, \text{ when } 0 < x < \pi$$

4

(c) Show that the rectified current through a half-wave rectifier is

$$I(t) = \frac{I_0}{\pi} - \frac{2I_0}{\pi} \left(\frac{1}{3} \cos 2\omega t + \frac{1}{15} \cos 4\omega t + \dots \right)$$

$$+ \frac{1}{2} I_0 \sin \omega t$$

4

(d) State and prove Parseval's theorem.

3

Or

Obtain the Fourier series for a triangular wave given by

$$y = 0 \quad \text{at } t = 0$$

$$y = a \quad \text{at } t = T/2$$

$$y = 0 \quad \text{at } t = T$$

3

www.prepnex.com